

The Application of GGE Biplot Analysis for Evaluating Test Locations and Mega-Environment Investigation of Cotton Regional Trials

[Favoris & Partage](#)[Export BibTex](#)[Export PDF](#)[Signaler](#)

Article de revue scientifique de Xu Naiying, Fok Michel, Zhang Guowei, Li Jian and Zhou Zhiguo - 2014

[Lien externe ou de téléchargement](#)

Référence bibliographique 2014, Xu Naiying, Fok Michel, Zhang Guowei, Li Jian and Zhou Zhiguo

The Application of GGE Biplot Analysis for Evaluating Test Locations and Mega-Environment Investigation of Cotton Regional Trials

Année de publication 2014

Auteur(s) Xu Naiying, Fok Michel, Zhang Guowei, Li Jian and Zhou Zhiguo

Titre du document The Application of GGE Biplot Analysis for Evaluating Test Locations and Mega-Environment Investigation of Cotton Regional Trials

Titre en anglais The Application of GGE Biplot Analysis for Evaluating Test Locations and Mega-Environment Investigation of Cotton Regional Trials

Adresse email de l'auteur naiyin@126.com

Adresse URL <file:///C:/Users/fok/Downloads/%EF%BC%88ASC-2013-0622%EF%BC%89The+Application+of+GGE+Biplot+Analysis+for+Evaluating+Test+Loc+Environment+Investigation+of+Cotton+Regional+Trials.pdf>

Journal ou Magazine Journal of Integrative Agriculture

Numéro du journal ou du magazine 9

Numéros de pages 1921-1933

Volume 13

Pays concerné(s) Chine

Thésaurus associé(s)

Enregistré le 2014-10-11

Modifié le 2014-10-11

Administré par [Fok Michel](#)

Résumé

In the process to the marketing of cultivars, identification of superior test locations within multi-environment variety trial schemes is of critical relevance. It is relevant to breeding organizations as well as to governmental organizations in charge of cultivar registration. Where competition among breeding companies exists, effective and fair multi-environment variety trials are of utmost importance to motivate investment in breeding. The objective of this study was to use genotype main effect plus genotype by environment interaction (GGE) biplot analysis to evaluate test locations in terms of discrimination ability, representativeness, and desirability, and to investigate the presence of multiple mega-environments in cotton production in the Yangtze River Valley, China. Four traits (cotton lint yield, fiber length, lint breaking tenacity, micronaire) and two composite selection indices were considered. It was found that the assumption of a single mega-environment in the Yangtze River valley for cotton production does not hold. The YaRV consists of three cotton mega-environments: a main one represented by 11 locations and two minor ones represented by two test locations each. This demands that the strategy of cotton variety registration or recommendation must be adjusted. GGE biplot analysis has also led to the identification of test location superior for cotton variety evaluation. Although test location desirable for selecting different traits varied greatly, Jinzhou was found to be desirable for selecting for all traits considered while Janyang was found to be desirable for none.

Résumé en anglais

In the process to the marketing of cultivars, identification of superior test locations within multi-environment variety trial schemes is of critical relevance. It is relevant to breeding organizations as well as to governmental organizations in charge of cultivar registration. Where competition among breeding companies exists, effective and fair multi-environment variety trials are of utmost importance to motivate investment in breeding. The objective of this study was to use genotype main effect plus genotype by environment interaction (GGE) biplot analysis to evaluate test locations in terms of discrimination ability, representativeness, and desirability, and to

investigate the presence of multiple mega-environments in cotton production in the Yangtze River Valley, China. Four traits (cotton lint yield, fiber length, lint breaking tenacity, micronaire) and two composite selection indices were considered. It was found that the assumption of a single mega-environment in the Yangtze River valley for cotton production does not hold. The YaRV consists of three cotton mega-environments: a main one represented by 11 locations and two minor ones represented by two test locations each. This demands that the strategy of cotton variety registration or recommendation must be adjusted. GGE biplot analysis has also led to the identification of test location superior for cotton variety evaluation. Although test location desirable for selecting different traits varied greatly, Jinzhou was found to be desirable for selecting for all traits considered while Janyang was found to be desirable for none.

[Aide](#) | [À propos de](#) | [Nous contacter](#) | [Informations légales](#)

[English](#) | [Français](#)

[AfomdNET](#) | [Cantool](#) | [CIRAD](#)

© Copyright CIRAD 2010-2020